

On Quantum Cohomology Rings of Fano Manifolds and a Formula of Vafa and Intriligator

Bernd Siebert¹ and Gang Tian

Courant Institute of Mathematical Sciences
251 Mercer Street, New York, NY 10012, USA

E-mail: siebert@cims.nyu.edu and tiang@cims.nyu.edu

Introduction

Quantum multiplications on the cohomology of symplectic manifolds were first proposed by the physicist Vafa [Va2] based on Witten's topological sigma models [Wil]. In [RuTi], Ruan and the second named author gave a mathematical construction of quantum multiplications on cohomology groups of *positive* symplectic manifolds (cf. Chapter 1). The construction uses the Gromov-Ruan-Witten invariants (GRW-invariants in the sequel) for semi-positive symplectic manifolds previously defined by Ruan [Ru]. A large class of such manifolds is provided by Fano manifolds (complex manifolds with ample anti-canonical bundle), including e.g. complex projective spaces, Del Pezzo surfaces and Grassmann manifolds. Let M be a Fano (or positive symplectic) manifold. The quantum cohomology

¹ On leave from: Mathematisches Institut, Bunsenstr. 3–5, 37073 Göttingen, Germany

$H_{[\omega]}^*(M)$ is just the cohomology space $H^*(M, \mathbb{C})$ with a (non-graded) associative, anti-commutative multiplication, the quantum multiplication. This multiplication depends on the choice of a (complexified) Kähler class $[\omega]$ on M . Its homogeneous part (the “weak coupling limit” $\lambda \cdot [\omega]$, $\lambda \rightarrow \infty$) is the usual cup product.

In this note we observe that quantum cohomology rings have a nice description in terms of generators and relations: If $H^*(M, \mathbb{C}) = \mathbb{C}[X_1, \dots, X_N]/(f_1, \dots, f_k)$ is a presentation of the cohomology ring (for simplicity we assume $\deg X_i$ even for the moment) then $H_{[\omega]}^*(M) = \mathbb{C}[X_1, \dots, X_N]/(f_1^{[\omega]}, \dots, f_k^{[\omega]})$, where $f_1^{[\omega]}, \dots, f_k^{[\omega]}$ are just the polynomials $f_1, \dots, f_k$ evaluated in the quantum ring associated to $[\omega]$ (Theorem 2.2). The $f_i^{[\omega]}$ actually are real-analytic in $[\omega]$ and thus have a natural analytic extension to $H^{1,1}(M)$, and the quantum cohomology rings fit together to form a flat analytic family over $H^{1,1}(M)$.

As an application of this observation we shall compute the quantum cohomology of the Grassmannians. The calculations for $G(k, n)$ reduce to the single quantum product $c_k \wedge_{\mathbb{Q}} s_{n-k}$ of the top non-vanishing Chern respectively Segre class of the tautological k -bundle. This turns out to be a simple exercise in linear algebra. We derive

Theorem 0.1 *Let S be the tautological bundle over the Grassmann manifold $G(k, n)$ of complex k -planes in $\mathbb{C}^n$, $[\omega] = -\lambda \cdot c_1(S)$ a $(1, 1)$ -class on $G(k, n)$ ($\lambda \in \mathbb{R}_{>0} \Leftrightarrow [\omega]$ Kähler). Then*

$$H_{[\omega]}^*(G(k, n)) = \mathbb{C}[X_1, \dots, X_k]/(Y_{n-k+1}, \dots, Y_{n-1}, Y_n + (-1)^k e^{-\lambda}),$$

where X_i corresponds to the i -th Chern class $c_i(S)$ and the Y_j (corresponding to the j -th Segre class of S) are given recursively by ($X_i = 0$ for $i > k$)

$$Y_j = -Y_{j-1} \cdot X_1 - \dots - Y_1 \cdot X_{j-1} - X_j.$$

This is in fact the form previously derived by Vafa using arguments from Quantum Field Theory [Va1], [Va2] (but note the sign of the quantum contribution). It also reduces the genus zero case of a nice conjectural formula for higher GRW-invariants of $G(k, n)$ stated in [Va1] and considered in a broader context in [In] to a purely algebraic problem. A mathematical formulation of the conjecture in terms of intersection theory on certain algebraic compactifications of moduli spaces of maps from a (fixed) Riemann surface to $G(k, n)$ was given by Bertram, Daskalopoulos and Wentworth [BDW]. They were able to verify the formula for genus one and $k = 2$.

This was the situation known to the authors around New Year 93/94. A preliminary version of this paper was spread in a limited number in the second half

of January 94. Since then some progress on related problems came to our knowledge: Witten (in a paper finished already in mid December 93) computed the quantum product $c_k \wedge s_{n-k}$, up to a check of genericity conditions. His arguments are more or less identical with our Lemma 3.3 (the content of Lemma 3.2 was taken for granted by him and should in fact be known classically). The underlying algebraic situation (the content of Chapter 2 and Chapter 4 of the present paper), however, and thus a proof of the structure of the quantum cohomology of $G(k, n)$ (even to a physical level of rigor), was left open. What nevertheless was inspiring for us, was his use of residues in the interpretation of coefficients of quantum products. The analysis of these residues lead to the first complete proof of the intriguing formula of Vafa and Intriligator for higher GRW-invariants of $G(k, n)$ (Theorem 4.6). On the way we found a nice (new?) interpretation of (Severi-Grothendieck-Griffiths) residues when there are “no components at infinity” (Proposition 4.1). An analogue of the formula of Vafa and Intriligator thus holds for all Fano manifolds whose cohomology ring is a commutative complete intersection, including the case where the corresponding sigma model has a Landau-Ginzburg description (Theorem 4.5). Later we realized that certain arguments in Chapter 4 are (encoded in physical language) already contained in the derivations of [Va1]. Moreover, some of the considerations in Chapter 2 have been found independently by Piunikhin [Pi].

Further partial support for the formula of Vafa/Intriligator were obtained in [RRW]. The quantum cohomology of (absolute) flag manifolds (containing $G(k, n)$ as special case) has been calculated in [GiKi] (complete flags) and [AsSa] (general case) *assuming* the existence of an equivariant version of quantum cohomology with certain functorial properties. After completion of this paper we received a preprint of Bertram containing the reduction to lower genus for the case of $G(k, n)$ [Be].

The first named author gratefully acknowledges support from the DFG (enabling him to visit the Courant Institute during the academic year 93/94) and he wishes to thank the Courant Institute for hospitality. The second author is partly supported by a NSF grant and an Alfred P. Sloan Fellowship.

1 Definition of quantum multiplications

In this section, we recall the definition of quantum multiplications given in [RuTi]. The definition uses the GRW-invariants as defined in [Ru].

A symplectic manifold (M, ω) of dimension $2n$ is called *(semi)-positive* if for any $R = f_*[S^2] \in H_2(M, \mathbb{Z})$, $f : S^2 \rightarrow M$, with $[\omega](R) > 0$ either $c_1(M)(R) > 0$ ($c_1(M)(R) \geq 0$). Any Fano manifold is positive w.r.t. any Kähler form. Let J

be a generic almost complex structure on M tamed by ω (i.e. $\omega(X, JX) > 0 \forall X \in TM \setminus \{0\}$). A J -holomorphic curve is a smooth map $f : \Sigma \rightarrow M$ satisfying $J \circ Df = Df \circ j$, where Σ is a Riemann surface (of genus g , say) and j is its standard complex structure. This last equation is a Cauchy-Riemann equation $\bar{\partial}f = 0$, $\bar{\partial} = \frac{1}{2}(D - J \circ D \circ j)$. For a technical reason it is convenient to look at the inhomogeneous equation $\bar{\partial}f = \gamma$ with γ (the pull-back to Σ of) a section of an appropriate bundle over $\Sigma \times M$. Solutions of this equation are called perturbed or (J, γ) -holomorphic curves. The GRW-invariant can be defined as follows:

Let $R \in H_2(M, \mathbb{Z})$ and $B_1, B_2, \dots, B_s$ be integral homology classes in $H_*(M, \mathbb{Z})$ satisfying:

$$\sum_{i=1}^s (2n - \deg B_i) = 2c_1(M)(R) + 2n(1 - g). \quad (\dim)$$

Every integral homology class can be represented by a so-called pseudo-manifold (a certain simplicial complex, cf. [RuTi]). For simplicity, we shall also use B_i to denote the pseudo-manifolds representing these homology classes. Then if (J, γ) is generic and B_i are in sufficiently general position (transversal w.r.t. the evaluation map $\Sigma \times \{f\} \rightarrow M$), for s generic points $t_1, \dots, t_s \in \Sigma$ there are only finitely many (J, γ) -holomorphic curves f satisfying: $f(t_i) \in B_i$, $i = 1, \dots, s$ and $f_*[\Sigma] = R$. We define $\tilde{\Phi}_{(R, \omega)}(B_1, \dots, B_s)$ to be the algebraic sum of such f with appropriate sign according to the orientation. One can prove that $\tilde{\Phi}_{(R, \omega)}(B_1, \dots, B_s)$ is independent of the choices of J , j , γ , and pseudo-manifolds representing $B_1, \dots, B_s$ provided the B_i are transversal to the Gromov boundary of the compactified moduli space of (J, γ) -holomorphic curves. The Gromov boundary here consists of curves $C = f(\Sigma) \cup g_1(S^2) \cup \dots \cup g_k(S^2)$ with: C connected; f a (J, γ) -holomorphic curve; g_ν a $(J, 0)$ -holomorphic rational curve (a “bubble”); $R = f_*[\Sigma] + \sum_\nu a_\nu g_{\nu*}[S^2]$ for some $a_\nu \in \mathbb{N}$. Transversality means that there are no such curves intersecting each B_i . Furthermore, $\tilde{\Phi}_{(R, \omega)}(B_1, \dots, B_s)$ depends only on the deformation class $\{\omega\}$ of ω . Therefore we obtain the GRW-invariant $\tilde{\Phi}_{(R, \{\omega\})}(B_1, \dots, B_s)$. As a matter of notation we define $\tilde{\Phi}_{(R, \omega)}(B_1, \dots, B_s)$ to be zero unless the dimensions match ($\dim$).

We now make a simple but important remark: Let J be an almost complex structure such that any J -holomorphic curve with $f_*[\Sigma] = R$ is regular, i.e. the linearization of the Cauchy-Riemann operator at f has trivial cokernel. Then $(J, 0)$ is generic and we can use exact J -holomorphic curves to compute the invariant $\tilde{\Phi}_{(R, \{\omega\})}(B_1, \dots, B_s)$. In case J is integrable and f is an immersion, regularity at f is equivalent to the vanishing of $H^1(C, N_{C|M})$, $C = f(\Sigma)$ [Gr, 2.1.B.]. In particular the invariant is easily computable in certain cases as follows:

Lemma 1.1 *Let (M, ω) be a Kähler manifold and $R \in H_2(M, \mathbb{Z})$ s.th. any holomorphic curve $C \subset M$ homologous to R is non-singular and has $H^1(C, N_{C|X}) = 0$.*

Let $B_1, \dots, B_s \subset M$ be complex submanifolds transversal to the evaluation map and the Gromov boundary. Then

$$\tilde{\Phi}_{(R, \{\omega\})}(B_1, \dots, B_s) = \sum_C \#(B_1 \cap C) \cdot \#(B_2 \cap C) \cdot \dots \cdot \#(B_s \cap C),$$

where the sum is over all holomorphic curves C homologous to R .

Note that $\#(B_1 \cap C) \cdot \dots \cdot \#(B_s \cap C)$ is precisely the number of ways to parametrize C by $f : \Sigma \rightarrow C$ with t_i mapping to B_i , $i = 1, \dots, s$. Moreover, there are no signs occurring in the formula since all spaces involved are canonically oriented by their complex structure.

Remark 1.2 (This will only be used to comment on the relationship of our work with [BDW].) More generally, if the moduli space $\mathcal{M}_{R, \Sigma}$ (for shortness $\mathcal{M}$ in the sequel) of holomorphic maps $f : \Sigma \rightarrow M$ with $f_*[\Sigma] = R$ has the expected dimension $c_1(M)(R) + n(1 - g)$ (equivalently, $H^1(\Sigma, f^*T_M) = 0$ for almost all f) the GRW-invariants $\tilde{\Phi}_R$ have an interpretation as intersection products of certain compactifications of $\mathcal{M}$: M being projective algebraic there clearly exist projective compactifications of $\mathcal{M}$. Choose one, say $\overline{\mathcal{M}}$. Then the evaluation map

$$\text{ev} : \Sigma \times \mathcal{M} \longrightarrow M, \quad (t, f) \longmapsto f(t)$$

extends rationally to $\overline{\mathcal{M}}$. Hence there is a modification $\pi : \Gamma \rightarrow \Sigma \times \overline{\mathcal{M}}$, biregular over $\Sigma \times \mathcal{M}$ such that ev has an extension $\tilde{\text{ev}} : \Gamma \rightarrow M$. For generic $t \in \Sigma$, $\Gamma_t := \pi^{-1}(\{t\} \times \mathcal{M})$ is a modification of $\overline{\mathcal{M}}$. We choose a common desingularization $\widehat{\mathcal{M}}$ of all the Γ_t , $t \in \Sigma$ generic (blow-up the sum of ideals $\mathcal{I}_t$ whose blow-up desingularizes Γ_t , then desingularize). Let $\iota_t : \widehat{\mathcal{M}} \rightarrow \Gamma_t \hookrightarrow \Gamma$. Then for $\beta_i \in H^*(M, \mathbb{Z})$ Poincaré-dual to B_i and generic choices of $t_1, \dots, t_s \in \Sigma$

$$\tilde{\Phi}_R(B_1, \dots, B_s) = (\iota_{t_1}^* \Phi^* \beta_1 \wedge \dots \wedge \iota_{t_s}^* \Phi^* \beta_s)[\widehat{\mathcal{M}}].$$

In fact, choosing $B_1, \dots, B_s$ as generic pseudo-manifolds then $(\Phi \circ \iota_{t_i})^{-1}(B_i)$ is Poincaré-dual to $\iota_{t_i}^* \Phi^* \beta_i$ and the bordism argument of [Ru] should generalize to show

$$\tilde{\Phi}_R(B_1, \dots, B_s) = \#(\Phi \circ \iota_{t_1})^{-1}(B_1) \cap \dots \cap (\Phi \circ \iota_{t_s})^{-1}(B_s),$$

($\#$ means algebraic sum with appropriate signs keeping track of the orientations). A proper proof has to provide a bordism to a generic situation carefully dealing with the singularities of $\mathcal{M}$ as well as with transversality and orientations. Details will be given in [RuTi]. $\diamond$

To define a quantum multiplication on $H^*(M, \mathbb{Z})$, we introduce the real-valued invariant

$$\tilde{\Phi}_{[\omega]}(B_1, \dots, B_s) = \sum_{R \in H_2(M, \mathbb{Z})} \tilde{\Phi}_{(R, \{\omega\})}(B_1, \dots, B_s) e^{-[\omega](R)}.$$

In general there might be infinitely many terms contributing to the sum (e.g. in the important and interesting case of Calabi-Yau manifolds) and one faces a non-trivial convergence problem. In the positive case, however, this sum is actually finite due to the dimension condition (dim). Let us assume M positive for the following (but see Remark 2.3). Note also that letting ω vary one gets back all the invariants $\tilde{\Phi}_{(R, \{\omega\})}(B_1, \dots, B_s)$ by solving some system of linear equations. The quantum multiplication $\wedge_Q$ on $H^*(M, \mathbb{R})$ is then characterized by the equation

$$(\alpha \wedge_Q \beta)[A] = \tilde{\Phi}_{[\omega]}(\alpha^\vee, \beta^\vee, A),$$

where $\alpha, \beta \in H^*(M, \mathbb{Z})$ and $^\vee$ means Poincaré-dual. In terms of a basis $\{A_i\}$ for the torsion free part of $H_*(M, \mathbb{Z})$ and $\{\alpha_i\}$ the Poincaré-dual basis of $H^*(M, \mathbb{Z})$ we may state this more explicitly as

$$\alpha_i \wedge_Q \alpha_j = \sum_{k,l} \eta^{lk} \tilde{\Phi}_{[\omega]}(A_i, A_j, A_k) \alpha_l,$$

with $(\eta^{lk})_{kl}$ inverse to the intersection matrix $(\eta_{ij})_{ij} = (A_i \cdot A_j)_{ij}$ ($\sum_k \eta_{ij} \eta^{kj} = \delta_{ik}$). The associativity of the quantum multiplication is highly non-trivial and shown by a careful analysis of the degeneration of rational curves (following an idea of Witten) in [RuTi].

A trivial, but decisive feature of this definition, that we are going to exploit, is that its homogeneous part reduces to the cup product

$$\alpha_i \wedge \alpha_j = \sum_{k,l} \eta^{lk} \cdot (A_i \cdot A_j \cdot A_k) \alpha_l.$$

2 A presentation for quantum cohomology

Denote by $\mathbb{C}\langle X_1, \dots, X_n \rangle$ the graded anticommutative $\mathbb{C}$ -algebra with generators X_i of degree d_i , i.e. with $X_i X_j = (-1)^{d_i d_j} X_j X_i$. If m of the X_i have odd degree this is isomorphic to $(\Lambda^* \mathbb{C}^m) \otimes (\text{Sym}^* \mathbb{C}^{n-m})$. We will call elements of this algebra ordered polynomials (since in addition to the coefficients one has to select an order among the factors in a monomial to determine its sign).

Let (M, ω) be a positive symplectic manifold and

$$H^*(M, \mathbb{C}) = \mathbb{C}\langle X_1, \dots, X_n \rangle / (f_1, \dots, f_k)$$

be a presentation of the cohomology ring, $f_i = \sum_{|J|=\deg f_i} a_{iJ} X^J$. (We use multiindex notation $J = (j_1, \dots, j_n)$, $X^J = X_1^{j_1} \wedge \dots \wedge X_n^{j_n}$, $|J| = \sum_{i=1}^n j_i d_i$ etc.) Denote by $\wedge_Q$ the product in the quantum cohomology $H_{[\omega]}^*(M)$. To distinguish clearly between calculations in $H^*(M, \mathbb{C})$ and in $H_{[\omega]}^*(M)$, we use a $\wedge$ to mark elements of the quantum cohomology ($\wedge$ might be thought of as a $\mathbb{C}$ -linear map $\mathbb{C}\langle X_1, \dots, X_n \rangle \rightarrow H_{[\omega]}^*(M)$). An ordered polynomial f may be evaluated on generators of the quantum cohomology provided the degrees match. We write $f(\hat{X}_1, \dots, \hat{X}_n)$ as in the commutative case. Especially, $\hat{X}^J := \hat{X}_1 \wedge_Q \dots \wedge_Q \hat{X}_1 \wedge_Q \dots \wedge_Q \hat{X}_n \wedge_Q \dots \wedge_Q \hat{X}_n$ with $\hat{X}_\nu$ occurring j_ν -often (note the difference from $(X^J)^\wedge$).

Lemma 2.1 $\hat{X}_1, \dots, \hat{X}_n$ generate $H_{[\omega]}^*(M)$.

Proof. By induction on the degree. So assume $\hat{X}_1, \dots, \hat{X}_n$ generate $H_{[\omega]}^*(M)$ up to degree $d - 1$. We want to show that the monomials $(X^J)^\wedge$, $|J| = d$, can be written as linear combination of (quantum) products of the $\hat{X}_\nu$. But by definition of the multiplication in $H_{[\omega]}^*(M)$

$$\hat{X}^J = (X^J)^\wedge + \sum_{|I| < d} a_I (X^I)^\wedge,$$

and by induction hypothesis $(X^I)^\wedge = \sum_{K \leq I} b_{IK} \hat{X}^K$, so

$$(X^J)^\wedge = \hat{X}^J - \sum_{\substack{K \leq I \\ |I| < d}} a_I b_{IK} \hat{X}^K.$$

◇

Next we are calculating $f_i(\hat{X}_1, \dots, \hat{X}_n)$ (i.e. in the quantum cohomology). Since f_i is a relation in the cohomology ring, the $(\deg f_i)$ -term vanishes. By the lemma we find an ordered polynomial $g_i^{[\omega]}$ (depending on $[\omega]$) in n variables (say $T_1, \dots, T_n$, $\deg T_i = \deg X_i$) of *lower degree* with

$$f_i(\hat{X}_1, \dots, \hat{X}_n) = g_i^{[\omega]}(\hat{X}_1, \dots, \hat{X}_n) \quad \text{in } H_{[\omega]}^*(M).$$

Thus $f_i^{[\omega]}(T_1, \dots, T_n) := f_i(T_1, \dots, T_n) - g_i^{[\omega]}(T_1, \dots, T_n) \in \mathbb{C}\langle T_1, \dots, T_n \rangle$ is a non-trivial relation between $\hat{X}_1, \dots, \hat{X}_n$.

Theorem 2.2 $H_{[\omega]}^*(M) = \mathbb{C}\langle T_1, \dots, T_n \rangle / (f_1^{[\omega]}, \dots, f_k^{[\omega]})$.

Proof. Let $\mathcal{J} \subset \mathbb{C}\langle T_1, \dots, T_n \rangle$ be the ideal of relations between the $\hat{X}_1, \dots, \hat{X}_n$. Then $(f_1^{[\omega]}, \dots, f_k^{[\omega]}) \subset \mathcal{J}$ and $H_{[\omega]}^*(M) = \mathbb{C}\langle T_1, \dots, T_n \rangle / \mathcal{J}$ by the lemma. Let

$F \in \mathcal{J} \setminus \{0\}$. Expand $F = F_d + F'$ with $F_d \neq 0$ (weighted) homogeneous of degree d , $d > 0$, and $\deg F' < d$. Then

$$F_d(\hat{X}_1, \dots, \hat{X}_n) = -F'(\hat{X}_1, \dots, \hat{X}_n)$$

in the quantum cohomology for F is a relation. But the highest degree ($\deg = d$) contribution to $F_d(\hat{X}_1, \dots, \hat{X}_n)$ is just $(F_d(X_1, \dots, X_n))^\wedge$. Its vanishing implies $F_d \in (f_1, \dots, f_k)$, i.e. $F_d = \varphi(f_1, \dots, f_k)$, φ a polynomial in k variables. Thus

$$\varphi(f_1^{[\omega]}, \dots, f_k^{[\omega]}) = F_d + F'' \quad \text{in } \mathbb{C}\langle T_1, \dots, T_n \rangle$$

with $\deg F'' < d$, and we may write $F = \varphi(f_1^{[\omega]}, \dots, f_k^{[\omega]}) + F' - F''$, $\deg F' - F'' < d = \deg F$. Proceeding by induction on the degree we finally see $F \in (f_1^{[\omega]}, \dots, f_k^{[\omega]})$, i.e. $\mathcal{J} = (f_1^{[\omega]}, \dots, f_k^{[\omega]})$ as claimed. $\diamond$

Remark 2.3 In the positive case $c_1(M) > 0$ the contributions with $R \neq 0$ give rise to terms of lower degree by the dimension condition, so we were able to fix $[\omega]$ and argue by considerations on the degree. Modulo convergence problems in the semi-positive case mentioned at the end of the previous chapter, Theorem 2.2 however remains valid. Arguments involving the degree can be replaced by the linear independence of terms $e^{-\lambda \cdot t}$ for various λ in the algebra $\mathbb{C}[[t]]$.

Another approach, which works without any positivity condition, is to use formal power series over $H_2(M, \mathbb{Z})$ (cf. also [Pi]): Let N be the (obvious) completion of the group ring over $H_2(M, \mathbb{Z})$. Then $H^*(M, \mathbb{Z}) \otimes_{\mathbb{Z}} N$ appears as natural domain of definition of quantum products for more general manifolds. Our arguments provide a presentation of the quantum cohomology ring as quotient of $\mathbb{Z}\langle T_1, \dots, T_n \rangle \otimes_{\mathbb{Z}} N$ by ordered polynomials with coefficients in N . Note also that in the positive case $H_{[\omega]}^*(M)$ may be viewed as homeomorphic image of this formal quantum cohomology ring by sending $R \in H_2(M, \mathbb{Z})$ to $e^{-[\omega](R)}$. $\diamond$

3 Grassmann manifolds

As is well-known (e.g. [Fu, Ex. 14.6.6]) the cohomology ring (in fact even the Chow ring) of the Grassmann variety $G(k, n)$ of k -planes in $\mathbb{C}^n$ has a presentation

$$H^*(G(k, n), \mathbb{C}) = \mathbb{C}[c_1, \dots, c_k] / (s_{n-k+1}, \dots, s_n),$$

with c_i corresponding to the Chern classes of the tautological k -bundle S and s_j to its Segre classes viewed as polynomials in $c_1, \dots, c_k$ via

$$\begin{aligned} (1 + c_1 + \dots + c_k)(1 + s_1 + s_2 + \dots) &= 1, \\ \text{i.e. } s_j &= -s_{j-1}c_1 - \dots - s_1c_{j-1} - c_j. \end{aligned} \tag{*}$$

Note that s_j is also the j -th Chern class of the universal quotient bundle Q , which has rank $n - k$, so $s_j = 0$ in $H^*(G(k, n), \mathbb{C})$ for $j > n - k$. In fact, as a polynomial in c_i , s_j lies in the relation ideal for $j > n$ by the recursion formula (*). Under the canonical isomorphism $\Phi : G(k, n) \simeq G(n - k, n)$, $\Lambda \mapsto (\mathbb{C}^n / \Lambda)^*$, S corresponds to the dual of the universal quotient bundle Q' on $G(n - k, n)$ and Q to the dual of the tautological $(n - k)$ -bundle S' . Thus c_i and s_j exchange (up to sign) their roles and one might as well write

$$H^*(G(k, n), \mathbb{C}) = \mathbb{C}[s_1, \dots, s_{n-k}] / (c_{k+1}, \dots, c_n),$$

this time with c_i polynomials in $s_1, \dots, s_{n-k}$ (this presentation is actually better adapted to Schubert calculus, i.e. geometry, see below). These remarks are made to emphasize the symmetry between c_i and s_j . Note also that the generators all have even degree, so we need not worry about questions of sign.

Schubert calculus (e.g. [Fu, § 14.7]) provides a basis of $H^*(G(k, n), \mathbb{C})$ as a $\mathbb{C}$ -vector space (indeed a basis of integral homology/cohomology as $\mathbb{Z}$ -module), indexed by tuples $(\lambda_1, \dots, \lambda_k)$, $n - k \geq \lambda_1 \geq \dots \geq \lambda_k \geq 0$, via

$$\{\lambda_1, \dots, \lambda_k\} := \det(s_{\lambda_i + j - i})_{1 \leq i, j \leq k},$$

i.e. by evaluating the Schur polynomial (S-function) associated to $(\lambda_1, \dots, \lambda_k)$ on the Segre classes of the tautological k -bundle ($s_j = 0$ for $j \notin \{0, \dots, n - k\}$). $\{\lambda_1, \dots, \lambda_k\}$ is (weighted) homogeneous of degree $2 \sum \lambda_i$. The Chern respectively Segre classes are given by

$$\begin{aligned} c_i &= (-1)^i \{1, \dots, 1, 0, \dots, 0\} \quad (\text{"1" } i\text{-times}), \\ s_j &= \{j, 0, \dots, 0\} \end{aligned}$$

(for c_i see [Fu, Lemma 14.5.1] or do an easy induction). The connection to classical Schubert calculus is given by Poincaré-duality. In fact, $\{\lambda_1, \dots, \lambda_k\} \cap [G(k, n)]$ may be represented by the Schubert varieties

$$\Omega_{\underline{V}}(n - k + 1 - \lambda_1, \dots, n - k + i - \lambda_i, \dots, n - \lambda_k).$$

To define the latter one has to fix a flag $\underline{V} = (V_1, \dots, V_n)$, $V_1 \subset \dots \subset V_n = \mathbb{C}^n$ of linear subspaces of $\mathbb{C}^n$, $\dim V_i = i$. Then for $(a_1, \dots, a_k)$, $0 \leq a_1 < \dots < a_k \leq n$

$$\Omega_{\underline{V}}(a_1, \dots, a_k) := \{\Lambda \in G(k, n) \mid \dim \Lambda \cap V_{a_i} \geq i, 1 \leq i \leq k\}.$$

The homology class $(a_1, \dots, a_k) := [\Omega_{\underline{V}}(a_1, \dots, a_k)]$ is independent of the choice of flag. We write $\{\underline{\lambda}\}^\vee := (n - k + 1 - \lambda_1, \dots, n - k + i - \lambda_i, \dots, n - \lambda_k) \in H_*(G(k, n), \mathbb{C})$. What is important for us is that $[G(k, n)]$ is Poincaré-dual to $\{0, \dots, 0\} = 1$ (trivial) and that the class of a point $[*]$ is Poincaré-dual to $\{n -$

$k, \dots, n-k\} = s_{n-k}^k = (-1)^{n-k} c_k^{n-k}$ (which makes sense as $\dim G(k, n) = k(n-k)$, but is less trivial: What is the Poincaré-dual of $c_1^{k(n-k)}$?). The intersection of classes of complementary dimension is especially easy: $|\underline{\lambda}| + |\underline{\mu}| = k(n-k)$, then (“duality theorem”):

$$\left(\{\underline{\lambda}\} \wedge \{\underline{\mu}\}\right) [G] = \{\underline{\lambda}\} \cap \{\underline{\mu}\}^\vee = \{\underline{\lambda}\}^\vee \cdot \{\underline{\mu}\}^\vee = \begin{cases} 1 & , \text{ if } \underline{\mu} = \underline{\lambda}^* \\ 0 & , \text{ otherwise.} \end{cases}$$

with $\underline{\lambda}^* := (n-k-\lambda_k, \dots, n-k-\lambda_1)$. In particular, we get $\eta_{\underline{\lambda}\underline{\mu}} = \delta_{\underline{\lambda}\underline{\mu}}$ for the intersection matrix. This ends our collection of facts concerning Grassmannians.

The product in $H_{[\omega]}^*(G(k, n))$ now reads

$$\widehat{\{\underline{\lambda}\}} \wedge_Q \widehat{\{\underline{\mu}\}} = \sum_{R \in H_2(G(k, n), \mathbb{Z})} \sum_{\underline{\nu}} \tilde{\Phi}_R \left(\{\underline{\lambda}\}^\vee, \{\underline{\mu}\}^\vee, \{\underline{\nu}\}^\vee \right) \widehat{\{\underline{\nu}^*\}} e^{-[\omega](R)}. \quad (**)$$

Next we observe that $H^{1,1}(G(k, n))$ is spanned by the single class $\{1, 0, \dots, 0\} = s_1 = -c_1$ and that $\det(Q) = (\det S)^*$ is actually very ample, the corresponding embedding being the Plücker embedding $\iota : G(k, n) \hookrightarrow \mathbb{P}(\Lambda^k \mathbb{C}^n)$ (so $s_1 = \iota^* c_1 \left(\mathcal{O}_{\mathbb{P}(\Lambda^k \mathbb{C}^n)}(1) \right)$). Dually, $H_2(G(k, n), \mathbb{Z})$ is spanned by the single class $\{n-k, \dots, n-k, n-k-1\}^\vee = (1, \dots, k-1, k+1) =: [L]$, and the sum over $H_2(G(k, n), \mathbb{Z})$ is just running through the set $\{d \cdot [L] \mid d \in \mathbb{N}_0\}$. Since $T_{G(k, n)} \simeq \text{Hom}(S, Q) \simeq S^* \otimes Q$, as one easily verifies by using the standard local coordinates on $G(k, n)$, $c_1(G(k, n)) = \text{rk}(Q) \cdot c_1(S^*) + \text{rk}(S^*) \cdot c_1(Q) = n \cdot s_1$.

Recall from Chapter 1 that $\tilde{\Phi}_{d \cdot [L]} \left(\{\underline{\lambda}\}^\vee, \{\underline{\mu}\}^\vee, \{\underline{\nu}\}^\vee \right) = 0$ unless $|\underline{\lambda}| + |\underline{\mu}| + |\underline{\nu}| = k(n-k) + c_1(G(k, n)) (d \cdot [L]) = k(n-k) + d \cdot n$. In calculating $s_{n-k+i}(\hat{c}_1, \dots, \hat{c}_k)$ (i.e. in $H_{[\omega]}^*(G(k, n))$), $\hat{c}_i$ the generators corresponding to c_i as in Chapter 2) we have $|\underline{\lambda}| + |\underline{\mu}| \leq n$, and “=” only in the case of s_n . But $|\underline{\nu}| \leq k(n-k) = \dim G(k, n)$, so we get no quantum contributions besides in case of s_n with $|\underline{\nu}| = k(n-k)$ and $d = 1$. Thus letting $\hat{\underline{c}} = (\hat{c}_1, \dots, \hat{c}_k)$, $s_{n-k+1}(\hat{\underline{c}}) = \dots = s_{n-1}(\hat{\underline{c}}) = 0$ in $H_{[\omega]}^*(G(k, n))$ and

$$\begin{aligned} s_n(\hat{\underline{c}}) &= -\hat{c}_1 s_{n-1}(\hat{\underline{c}}) - \dots - \hat{c}_{k-1} s_{n-k+1}(\hat{\underline{c}}) - \hat{c}_k s_{n-k}(\hat{\underline{c}}) \\ &= -\hat{c}_k s_{n-k}(\hat{\underline{c}}) = -\tilde{\Phi}_{[L]} \left(c_k^\vee, s_{n-k}^\vee, [*] \right) \cdot e^{-[\omega](L)}, \end{aligned}$$

where for the last equality we have used (**). Taking into account Theorem 2.2 to prove Theorem 0.1 we are left with

Proposition 3.1 $\tilde{\Phi}_{[L]} \left(c_k^\vee, s_{n-k}^\vee, [*] \right) = (-1)^k$.

For the proposition we are first classifying holomorphic curves homologous to $[L] = (1, \dots, k-1, k+1)$.

Lemma 3.2 *Let $C \subset G(k, n)$ be a (rational) curve homologous to $(1, \dots, k-1, k+1)$. Then C is a Schubert variety $\Omega_{\underline{V}}(1, \dots, k-1, k+1)$, i.e. there are linear subspaces $U \subset W \subset \mathbb{C}^n$, $\dim U = k-1$, $\dim W = k+1$, with $C = \{\Lambda \in G(k, n) \mid U \subset \Lambda \subset W\}$.*

Proof. Since $s_1[C] = 1$ by the duality theorem, $\deg \iota(C) = 1$, so the image of C under the Plücker embedding $\iota : G(k, n) \rightarrow \mathbb{P}(\Lambda^k \mathbb{C}^n)$ is a linear $\mathbb{P}^1$. The image of $G(k, n)$ consists precisely of (rays of) *decomposable* vectors $v_1 \wedge \dots \wedge v_k = \iota(\langle v_1, \dots, v_k \rangle) \in \Lambda^k \mathbb{C}^n$. Locally the vectors $v_1, \dots, v_k$ may be chosen to vary smoothly with $\Lambda \in G(k, n)$: In fact, the standard (affine) coordinate neighbourhood of $\Lambda \in G(k, n)$ is $\text{Hom}(\Lambda, \mathbb{C}^n/\Lambda)$ with $\Psi : \text{Hom}(\Lambda, \mathbb{C}^n/\Lambda) \rightarrow G(k, n)$, $\varphi \mapsto \langle v + \varphi(v) \mid v \in \Lambda \rangle$ (in particular $0 \in \text{Hom}(\Lambda, \mathbb{C}^n/\Lambda)$ corresponds to Λ). Now fixing a basis $v_1, \dots, v_k$ of Λ , $\iota \circ \Psi$ may be represented (lifted to $\Lambda^k \mathbb{C}^n$) by

$$\varphi \mapsto (v_1 + \varphi(v_1)) \wedge \dots \wedge (v_k + \varphi(v_k)).$$

Thus choosing $\Lambda' \in C$ sufficiently close to $\Lambda \in C$ and letting $e_1, \dots, e_l$ be a basis of $\Lambda \cap \Lambda'$, completed by $e_{l+1}, \dots, e_k$ and $e'_{l+1}, \dots, e'_k$ to a basis of Λ and Λ' respectively, we have for $t \in \mathbb{C}$ small

$$e_1 \wedge \dots \wedge e_l \wedge e_{l+1} \wedge \dots \wedge e_k + t \cdot e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = v_1(t) \wedge \dots \wedge v_k(t)$$

with $v_i(0) = e_i$ by construction. Taking $\left. \frac{d}{dt} \right|_{t=0}$ yields

$$e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = \dot{v}_1(0) \wedge e_2 \wedge \dots \wedge e_k + \dots + e_1 \wedge \dots \wedge e_{k-1} \wedge \dot{v}_k(0).$$

As one sees by expanding $\dot{v}_i(0)$ in terms of a basis of $\mathbb{C}^n$ containing $\{e_1, \dots, e_k, e'_{l+1}, \dots, e'_k\}$ we may gather the linearly independent terms with e_k and without e_k to form two equations. The left-hand side of the equation above belongs to the latter ($\Lambda \neq \Lambda' \Rightarrow l < k$), so we get

$$e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = e_1 \wedge \dots \wedge e_{k-1} \wedge (\dot{v}_k(0) - \lambda \cdot e_k)$$

for some $\lambda \in \mathbb{C}$ (s.th. $\dot{v}_k(0) - \lambda \cdot e_k$ lies in the span of the basis vectors different from e_k). By linear independence of wedge products of a basis of $\mathbb{C}^n$ this shows $l = k-1$. In view of the linearity of $\iota(C)$ we conclude

$$\iota(C) = \left\{ [t \cdot e_1 \wedge \dots \wedge e_{k-1} \wedge e'_k + u \cdot e_1 \wedge \dots \wedge e_{k-1} \wedge e_k] \mid [t : u] \in \mathbb{P}^1 \right\},$$

so C is the Schubert variety $\Omega_{\underline{V}}(1, \dots, k-1, k+1)$ belonging to a flag $\underline{V}$ with $V_{k-1} = \langle e_1, \dots, e_{k-1} \rangle =: U$ and $V_{k+1} = \langle e_1, \dots, e_{k-1}, e_k, e'_k \rangle =: W$. $\diamond$

Lemma 3.3 *Let $A_1 = \Omega_{\underline{V}^1}(n-k, n-k+1, \dots, n-1) = \{1, \dots, 1\}^\vee = (-1)^k c_k^\vee$, $A_2 = \Omega_{\underline{V}^2}(1, n-k+2, \dots, n) = \{n-k, 0, \dots, 0\}^\vee = s_{n-k}^\vee$, $A_3 = \{*\} = \Omega_{\underline{V}^3}(1, \dots, k) = \{n-k, \dots, n-k\}^\vee$, where $\underline{V}^1, \underline{V}^2, \underline{V}^3$ are three transversal flags (i.e. $\dim V_i^1 \cap V_j^2 \cap V_k^3 = \max\{0, i+j+k-2n\}$). Then there is one and only one rational curve C homologous to $[L]$ and with $C \cap A_i \neq \emptyset$, $i = 1, 2, 3$. Moreover, $C \cdot A_1 = C \cdot A_2 = C \cdot A_3 = 1$.*

The lemma can be proved by doing intersection theory on the flag manifold $F(k-1, k+1; n)$, which parametrizes rational curves of minimal degree by the preceding lemma, and using the two obvious maps $\pi : F(k-1, k+1; n) \rightarrow G(k, n)$ and $p : F(k-1, k+1; n) \rightarrow G(k, n)$ (calculate $(p_*\pi^*[A_1]) \cdot (p_*\pi^*[A_2]) \cdot (p_*\pi^*[A_3])$). This method might be appropriate for more general A_1, A_2, A_3 , yet in our case an explicit linear algebra argument is simpler and even more enlightening.

Proof. We need to find three k -planes $\Lambda^1, \Lambda^2, \Lambda^3$ and subspaces $U, W \subset \mathbb{C}^n$, $\dim U = k-1$, $\dim W = k+1$ with

1. $U \subset \Lambda^1 \cap \Lambda^2 \cap \Lambda^3 \subset \Lambda^1 + \Lambda^2 + \Lambda^3 \subset W$.
2. $\dim \Lambda^1 \cap V_{n-k+(i-1)}^1 \geq i$, $i = 1, \dots, k$.
3. $V_1^2 \subset \Lambda^2$.
4. $\Lambda^3 = V_k^3$,

(1) says that $\Lambda^1, \Lambda^2, \Lambda^3$ lie on the rational curve C defined by U and W according to Lemma 3.2, whereas (2)–(4) rephrase the conditions $\Lambda_i = A_i \cap C$, $i = 1, 2, 3$ respectively. We are now using transversality of the flags $\underline{V}^1, \underline{V}^2, \underline{V}^3$. From (3), (4) and (1) we readily deduce $W = V_k^3 + V_1^2$, and (2) with $i = k$ shows $\Lambda^1 \subset V_{n-1}^1$, so by (4) and (1) we get $U = V_k^3 \cap V_{n-1}^1$. This choice of U, W implies $\Lambda^1 = W \cap V_{n-1}^1$, $\Lambda^2 = U + V_1^2$, $\Lambda^3 = V_k^3$. Conversely these $\Lambda^1, \Lambda^2, \Lambda^3$ fulfill (1)–(4). $\diamond$

Proof of Proposition 3.1. In view of the preceding lemma and the criterion for multiplicity one (Lemma 1.1 with $R = [L]$) the only thing remaining to be checked is transversality of A_i w.r.t. the evaluation map

$$\Sigma \times \{f : \Sigma \rightarrow G(k, n)\} \rightarrow G(k, n).$$

The moduli space $\{f\}$ of holomorphic curves in question is in our case isomorphic to the flag manifold $F(k-1, k+1; n) = \{(U, W)\}$ by Lemma 3.2 and 3.3. But $\Lambda_i = A_i \cap f(\Sigma)$ clearly varies when f (and hence (U, W)) varies as seen explicitly

in the proof of Lemma 3.3. Note also that the Gromov boundary is empty in this case. This follows either from the explicit description of the moduli space as flag manifold or from the fact that $L \in H_2(G(k, n), \mathbb{Z})$ is primitive and thus can not be represented by any reducible holomorphic curve. $\diamond$

This finishes the proof of Theorem 0.1.

4 Higher invariants

The main results of [RuTi] show how to compute higher GRW-invariants (i.e. with more than three entries or for higher genus Riemann surfaces) from the genus 0 three-point functions inductively. Namely, for $g > 0$

$$\tilde{\Phi}_{[\omega]}^g(B_1, \dots, B_s) = \sum_{i,j} \eta^{ij} \tilde{\Phi}_{[\omega]}^{g-1}(B_1, \dots, B_s, A_i, A_j),$$

(η_{ij}) the intersection matrix with respect to a basis $\{A_i\}$ of $H^*(M, \mathbb{C})$. More invariantly, the right-hand side of course is the trace with respect to η of the bilinear form

$$H^*(M, \mathbb{R}) \times H^*(M, \mathbb{R}) \longrightarrow \mathbb{R}, \quad (B', B'') \longmapsto \tilde{\Phi}_{[\omega]}^{g-1}(B_1, \dots, B_s, B', B'').$$

Secondly, for $g = 0$ and $1 < r < s - 1$ (otherwise trivial) we have the *composition law*

$$\tilde{\Phi}_{[\omega]}^0(B_1, \dots, B_s) = \sum_{i,j} \eta^{ij} \tilde{\Phi}_{[\omega]}^0(B_1, \dots, B_r, A_i) \tilde{\Phi}_{[\omega]}^0(A_j, B_{r+1}, \dots, B_s),$$

a trace with respect to η as well (for $s = 4$ and $r = 2$ this equation states the associativity of quantum products.) Our goal in this section is to give a closed formula for higher invariants in terms of the relations $f_1^{[\omega]}, \dots, f_k^{[\omega]}$ of the quantum cohomology ring. Putting all $\tilde{\Phi}_{[\omega]}^g$ together for different s we get a $\mathbb{C}$ -linear map

$$\langle \rangle_g : \mathbb{C}\langle X_1, \dots, X_n \rangle \longrightarrow \mathbb{C}, \quad X_1^{\nu_1} \dots X_n^{\nu_n} \longmapsto \tilde{\Phi}_{[\omega]}^g(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, \dots, X_n^\vee),$$

the Poincaré-dual $X_i^\vee$ of X_i occurring ν_i -times. But from the composition law denoting by α_i the Poincaré-dual of A_i

$$\hat{X}_1^{\nu_1} \wedge_Q \dots \wedge_Q \hat{X}_n^{\nu_n} = \sum_{ij} \eta^{ij} \tilde{\Phi}_{[\omega]}^0(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, A_i) \hat{\alpha}_j,$$

so $\langle X_1^{\nu_1} \dots X_n^{\nu_n} \rangle_0 = \tilde{\Phi}_{[\omega]}^0(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, [M])$ is nothing but the coefficient of the class $[\Omega]$ of the normalized volume form in $\hat{X}_1^{\nu_1} \wedge_Q \dots \wedge_Q \hat{X}_n^{\nu_n}$. That is, $\langle \rangle_0$ decomposes

$$\mathbb{C}\langle X_1, \dots, X_n \rangle \longrightarrow \mathbb{C}\langle X_1, \dots, X_n \rangle / (f_1^{[\omega]}, \dots, f_k^{[\omega]}) \simeq H_{[\omega]}^*(M) \xrightarrow{\text{top}} H^{2n}(M, \mathbb{C}),$$

and $H^{2n}(M, \mathbb{C})$ is identified with $\mathbb{C}$ by sending $[\Omega]$ to 1. In case $n = k$ and the generators have even degree, i.e. $H^*(M, \mathbb{C})$ a (commutative) complete intersection ring, one can use higher dimensional residues to express this map more explicitly (we refer the reader to [GrHa] and [Ts] for the general facts on residues to be used):

Recall that the residue of $F \in \mathbb{C}[X_1, \dots, X_k]$ with respect to a polynomial mapping $g = (g_1, \dots, g_k) : \mathbb{C}^k \rightarrow \mathbb{C}^k$ with $g^{-1}(0)$ finite (or equivalently, $\mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k)$ is artinian, i.e. finite dimensional as vector space over $\mathbb{C}$) in $a \in g^{-1}(0)$ is defined by

$$\text{res}_g(a; F) := \frac{1}{(2\pi i)^k} \int_{\Gamma_a^\varepsilon} \frac{F}{g_1 \cdots g_k} dX_1 \cdots dX_k,$$

with $\Gamma_a^\varepsilon = \{x \in U(a) \mid |g_i(x)| = \varepsilon\}$, $U(a)$ a neighbourhood of a with $g^{-1}(0) \cap U(a) = \{a\}$ and ε so small that Γ_a^ε lies relatively compact in $U(a)$. Γ_a^ε is smooth for almost all ε by Sard's Theorem and has a canonical orientation by the k -form $d(\arg g_1) \wedge \dots \wedge d(\arg g_k)|_{\Gamma_a^\varepsilon}$. (This local residue of course makes sense for holomorphic g and $F \in \mathcal{O}_a$, but the polynomial case, to which the general case may easily be reduced, is sufficient for our purposes). We define the total residue

$$\text{Res}_g(F) := \sum_{a \in g^{-1}(0)} \text{res}_g(a; F),$$

which is also known as Grothendieck residue symbol $\left(\frac{F}{g_1, \dots, g_k}\right)$ in the context of duality theory in algebraic geometry [Ha]. Let $J = \det\left(\frac{\partial g_i}{\partial X_j}\right)$ be the Jacobian of g . Then for regular values y of g

$$\text{Res}_{g-y}(F) = \sum_{x \in g^{-1}(y)} \left(\frac{F}{J}\right)(x) = \text{tr}\left(\frac{F}{J}\right)(y).$$

Therefore $\text{tr}(F/J)$ extends holomorphically (surprise!) to a neighbourhood of 0 (the extension will be denoted $\text{tr}(F/J)$ as well) and

$$\text{Res}_g(F) = \text{tr}\left(\frac{F}{J}\right)(0).$$

One abstract feature in our setting is that we have weights d_i associated to X_i and that our relations $f_i^{[\omega]}$ form a standard basis of the relation ideal with respect to these weights, i.e.

$$(\text{In } f_1^{[\omega]}, \dots, \text{In } f_k^{[\omega]}) = (f_1, \dots, f_k) = \text{In}(f_1^{[\omega]}, \dots, f_k^{[\omega]}),$$

where “In” means taking initial forms. This is trivial in our case since we started with the homogeneous generators $f_1, \dots, f_k$ of the relation ideal in a presentation of $H^*(M, \mathbb{C})$. In this situation one can describe the residue map algebraically as follows:

Proposition 4.1 *Let $R = \mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k)$ be artinian such that $\{g_i\}$ is a standard basis of the relation ideal with respect to weights d_i of X_i . Put $N := \sum_i \deg g_i - \sum_i d_i$, $R_{<N} := \{F \in \mathbb{C}[X_1, \dots, X_k] \mid \deg F < N\}/(g_1, \dots, g_k)$ and $J = \det\left(\frac{\partial g_i}{\partial X_j}\right)$. Then*

$$R = R_{<N} \oplus \mathbb{C} \cdot J,$$

and the total residue map $\text{Res}_g : \mathbb{C}[X_1, \dots, X_k] \rightarrow \mathbb{C}$ factorizes via the projection onto the second factor as follows

$$\mathbb{C}[X_1, \dots, X_k] \xrightarrow{\text{can}} R \xrightarrow{\text{pr}_2} \mathbb{C} \cdot J \longrightarrow \mathbb{C},$$

where the last map sends J to $\dim_{\mathbb{C}} R$.

Proof. To check the normalization we observe $\text{Res}_g(J) = \text{tr}_g(1)(0) = \text{degree of } g \text{ over } 0 = \dim_{\mathbb{C}} R$ (the latter equality is generally true by flatness if the covering space is Cohen-Macaulay [Fi]). Next it is well known that the residue vanishes on elements of $(g_1, \dots, g_k)$. The claim thus reduces to $\ker(\text{Res}_g)/(g_1, \dots, g_k) = R_{<N}$.

If $\deg F < N$ then $F dX_1 \dots dX_k / g_1 \dots g_k$ extends to a rational differential form φ on weighted projective space $V = \mathbb{P}_{(1, d_1, \dots, d_k)}$ with polar divisor $D_1 + \dots + D_k \in \text{Div}(V)$, D_i the natural extension of the divisor (g_i) to V . The point of course is that the divisor at infinity $V \setminus \mathbb{C}^k$ is not a polar divisor of φ . We claim $|D_1| \cap \dots \cap |D_k| \subset \mathbb{C}^k$. In fact, the restriction of the homogenization of g_i to $V \setminus \mathbb{C}^k \simeq \mathbb{P}_{(d_1, \dots, d_k)}$ is just $\text{In } g_i$ and $V(\text{In } g_1, \dots, \text{In } g_k) = \{0\} \in \mathbb{C}^k$. The latter follows because otherwise $\dim \text{Spec } \mathbb{C}[X_1, \dots, X_k]/(\text{In } g_1, \dots, \text{In } g_k) > 0$ by homogeneity. But from the standard basis property we have $\dim_{\mathbb{C}} \mathbb{C}[X_1, \dots, X_k]/(\text{In } g_1, \dots, \text{In } g_k) = \dim_{\mathbb{C}} \mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k) < \infty$. — We may thus desingularize V (at infinity) without violating the discreteness of $|D_1| \cap \dots \cap |D_k|$ (we use the same notations for the pulled-back objects). Now the global residue theorem tells that on the compact manifold V the sum of the local residues of φ with respect to $D_1, \dots, D_k$ equals zero (the local residue $\text{res}_g(a; F)$ is coordinate free in so far that it depends only on the associated rational differential form $F dX_1 \dots dX_k / g_1 \dots g_k$ and the divisors $(g_1), \dots, (g_k)$). This proves $\text{Res}_g(F) = 0$ in case $\deg F < N$.

The second case is F homogeneous of degree $> N$. We show $F \in (\text{In } g_1, \dots, \text{In } g_k)$. This is an easy generalization of a theorem of Macaulay (cf. e.g. [Ts]) to the weighted situation. Namely, for any G (wlog. homogeneous), setting $P = F \cdot G$, $Q = \text{In}(g_1) \dots \text{In}(g_k)$, we have $\frac{P}{Q} dX_1 \dots dX_k = (\deg P - \deg Q + \sum_i d_i)^{-1} d\sigma = (\deg P - N)^{-1} d\sigma$ with

$$\sigma = \frac{P}{Q} \sum_{j=1}^k (-1)^{j-1} d_j \cdot X_j dX_1 \dots \widehat{dX_j} \dots dX_k,$$

where $\frown$ means that this entry is to be left out. This is a simple check using the weighted Euler formula $\sum_j d_j X_j \frac{\partial H}{\partial X_j} = \deg(H) \cdot P$, H weighted homogeneous (same proof as usual). Thus

$$\text{Res}_{\text{In}g}(F \cdot G) = \sum_a \int_{\Gamma_a^\varepsilon} d\sigma = 0$$

for all (homogeneous) $G \in \mathbb{C}[X_1, \dots, X_k]$. But this implies $F \in (\text{In}g_1, \dots, \text{In}g_k)$ (“duality theorem”, cf. [Ts] — this is Poincaré duality in our case!). Modulo $(g_1, \dots, g_k)$ this means that we may reduce F to lower degree. So proceeding by induction N turns out to be “top-degree” in R in that all elements of R can be represented by polynomials of degree $\leq N$.

What remains to be checked is that for F homogeneous of degree N , $\text{Res}_g(F) \neq 0$ or $F/(g_1, \dots, g_k) \in R_{<N}$. But in the first instance $\text{Res}_{\text{In}(g)}(F \cdot G) = 0 \forall G \in \mathbb{C}[X_1, \dots, X_k]$ as shown above, and again $F \in (\text{In}g_1, \dots, \text{In}g_k)$, i.e. modulo $(g_1, \dots, g_k)$, F may be represented by a polynomial of degree $< N$. $\diamond$

Remark 4.2 The decomposition $R = R_{<N} \oplus \langle J \rangle$ is *not* canonical but rather depends on the particular presentation of R . This may be seen either in elementary terms from the transformation formula for residues or as manifestation of the choice of an isomorphism $\text{Ext}_V^k(\mathcal{O}_Z, \Omega_V^k) \simeq H^0(V, \mathcal{O}_Z)$ in the duality morphism

$$\text{Ext}_V^k(\mathcal{O}_Z, \Omega_V^k) \times H^0(V, \mathcal{O}_Z) \longrightarrow \mathbb{C}$$

induced by the global residue. For quantum cohomology rings and weightings coming from cohomology, however, $R_{<N} = \oplus_{d < N} H^d(M, \mathbb{C})$, $N = \dim_{\mathbb{C}} M$ and $\langle Y \rangle = H^{2N}(M, \mathbb{C})$, so the decomposition *has* an invariant meaning in this case. $\diamond$

In the quantum cohomology ring the top-degree class $\mathbb{C} \cdot J$ is thus spanned by the class $[\Omega]$ of the volume form. Let $F_{[\Omega]}$ be a polynomial of degree N representing $[\Omega]$ (modulo $(f_i^{[\omega]})$ or modulo (f_i) , this will yield the same result), and set $c = 1/\text{Res}_{f^{[\omega]}}(F_{[\Omega]})$. By the interpretation of $\langle F \rangle_0$ as coefficient of $[\Omega]$ of $F(\hat{X}_1, \dots, \hat{X}_k) \in H_{[\omega]}^*(M)$ we conclude (for $H^*(M, \mathbb{C})$ a complete intersection)

$$\langle F \rangle_0 = c \cdot \text{Res}_{f^{[\omega]}}(F) = c \cdot \text{tr}_{f^{[\omega]}} \left(\frac{F}{J} \right) (0).$$

To incorporate the higher genus case we prove

Lemma 4.3 *Notations as in the proposition and $F \in \mathbb{C}[X_1, \dots, X_k]$ let B_F be the bilinear form $R \times R \rightarrow \mathbb{C}$, $(\alpha, \beta) \mapsto \text{Res}_g(F \cdot G_\alpha \cdot G_\beta)$ (with $G_\alpha, G_\beta \in \mathbb{C}[X_1, \dots, X_k]$ representing α, β — this is well-defined), $\eta = B_1$ (“intersection form”). Then*

$$\text{tr}_\eta B_F = \text{Res}_g(F \cdot J).$$

Proof. We give a basis-free, algebraic proof (some might find the brute force method by adapting a basis to η more enlightening).

By definition $\text{tr}_\eta B_F$ is the trace of the endomorphism $\mu_F : R \rightarrow R$ of multiplication by (the class of) F . Putting $Z = \text{Spec } R$ we have $R = \bigoplus_{z \in g^{-1}(0)} \mathcal{O}_{Z,z}$. For $z \in g^{-1}(0)$, $F - F(z)$ is nilpotent in $\mathcal{O}_{Z,z}$ (for it has value 0 in z), so $\mu_{F-F(z)}$ has trace 0 and

$$\text{tr}(\mu_F | \mathcal{O}_{Z,z}) = \text{tr}(\mu_{F-F(z)} | \mathcal{O}_{Z,z}) + \text{tr}(\mu_{F(z)} | \mathcal{O}_{Z,z}) = F(z) \cdot \dim_{\mathbb{C}} \mathcal{O}_{Z,z}.$$

Furthermore, $\dim_{\mathbb{C}} \mathcal{O}_{Z,z} = \deg_z(g)$, the local mapping degree of g at z , so summing up we get

$$\text{tr}_\eta B_F = \sum_{z \in g^{-1}(0)} \deg_z(g) \cdot F(z) = \text{tr}_g(F),$$

which is nothing but $\text{Res}_g(F \cdot J)$ as claimed. $\diamond$

In view of the reduction formula to lower genus stated above we conclude a rigorous version of the “handle gluing formula”, previously established in QFT by Witten [Wi2].

Proposition 4.4 *Let $H^*(M, \mathbb{C}) = \mathbb{C}[X_1, \dots, X_k] / (f_1, \dots, f_k)$ (the commutative complete intersection case) and $f_i^{[\omega]}$ the induced relations in $H_{[\omega]}^*(M)$ as in Theorem 2.2. Put $J = \det \left(\frac{\partial f_i^{[\omega]}}{\partial X_j} \right)$. Then*

$$\langle F \rangle_g = \langle J \cdot F \rangle_{g-1}$$

holds for all $F \in \mathbb{C}[X_1, \dots, X_k]$. $\diamond$

In other words, multiplication by J acts as “attaching a handle” to our Riemann surface. Our considerations so far yield the main result of the present chapter:

Theorem 4.5 *Assumptions as in the preceding proposition then for all $F \in \mathbb{C}[X_1, \dots, X_k]$ the following holds*

$$\langle F \rangle_g = c \cdot \text{Res}_{f^{[\omega]}}(J^g \cdot F) = c \cdot \lim_{\substack{y \rightarrow 0 \\ y \text{ regular} \\ \text{value of } f^{[\omega]}}} \text{tr}_{f^{[\omega]}}(J^{g-1} \cdot F)(y),$$

with $c = 1/\text{Res}_{f^{[\omega]}}(F_{[\Omega]})$, $F_{[\Omega]}$ a polynomial representing the class $[\Omega] \in H_{[\omega]}^*(M)$ of the normalized volume form. $\diamond$

We emphasize that for $g > 0$ or $0 \in \mathbb{C}^k$ a regular value of $f^{[\omega]}$ the right-hand side has the form $\sum_{\nu} a_{\nu} F(y_{\nu})$ with $g^{-1}(0) = \{y_{\nu}\}$ and constants a_{ν} independent of F !

Note also that 0 is a regular value of $f^{[\omega]}$ iff $\dim_{\mathbb{C}} \mathcal{O}_{Z,z} = 1$ for all $z \in (f^{[\omega]})^{-1}(0)$, $Z = \text{Spec } \mathbb{C}[X_1, \dots, X_k]/(f_1^{[\omega]}, \dots, f_k^{[\omega]})$, which is if and only if

$$\dim_{\mathbb{C}} H^*(M, \mathbb{C}) = \sharp \left((f^{[\omega]})^{-1}(0) \right)$$

(“ $\geq$ ” always).

For the case of $G(k, n)$ we have a basis of Schubert classes $\{\lambda_1, \dots, \lambda_k\}$ parametrized by sequences $n - k \geq \lambda_1 \geq \dots \geq \lambda_k \geq 0$. The latter are in 1–1 correspondence with subsets $\{\lambda_k + 1, \lambda_{k-1} + 2, \dots, \lambda_1 + k\}$ of $\{1, \dots, n\}$, so $\dim_{\mathbb{C}} H^*(G(k, n), \mathbb{C}) = \binom{n}{k}$.

To find $\binom{n}{k}$ distinct elements in $(f^{[\omega]})^{-1}(0)$ we use a description of the cohomology ring coming from the study of the corresponding Landau-Ginzburg model in physics [Va1]: There is a function W (“Landau-Ginzburg potential”) s.th. in the notations of Theorem 0.1

$$Y_{n+1-i}(X_1, \dots, X_k) = \frac{\partial W}{\partial X_i},$$

and then $f_i^{[\omega]} = \frac{\partial W^{[\omega]}}{\partial X_i}$ with $W^{[\omega]} := W + (-1)^k e^{-[\omega](L)} \cdot X_1$. W has a simple description in terms of Chern roots, i.e. after composition with $\Sigma : \mathbb{C}^k \rightarrow \mathbb{C}^k$, $(\underline{\lambda}) = (\lambda_1, \dots, \lambda_k) \mapsto \Sigma(\underline{\lambda}) = (-\sigma_1(\underline{\lambda}), \sigma_2(\underline{\lambda}), \dots, (-1)^k \sigma_k(\underline{\lambda}))$, σ_i the elementary symmetric polynomials: $W \circ \Sigma = -\frac{1}{n+1} \sum_i \lambda_i^{n+1}$. Thus

$$W^{[\omega]} \circ \Sigma = - \sum_{i=1}^k \left(\frac{\lambda_i^{n+1}}{n+1} - (-1)^k e^{-[\omega](L)} \cdot \lambda_i \right).$$

This all is an easy formal consequence of the algebraic relations between Chern and Segre classes on one side and Chern roots on the other, cf. [BDW] for a mathematical account. Now $\frac{\partial W^{[\omega]}}{\partial X_i}(\Sigma(\underline{\lambda})) = 0$, $i = 1, \dots, k$, if $\frac{\partial W^{[\omega]} \circ \Sigma}{\partial \lambda_i}(\underline{\lambda}) = 0$ and if Σ is non-degenerate in $(\underline{\lambda})$. The latter obviously is equivalent to $\lambda_i \neq \lambda_j$ for $i \neq j$. Solving the second equation means

$$\lambda_i^n = (-1)^k e^{-[\omega](L)}, \quad i = 1, \dots, k$$

which has $n \cdot (n-1) \cdots (n-k+1)$ solutions with distinct λ_i . The fiber of Σ over the regular value $\Sigma(\underline{\lambda})$ consists of the $k!$ permutations of $\{\lambda_1, \dots, \lambda_k\}$, so we get precisely $\binom{n}{k}$ distinct elements of $(f^{[\omega]})^{-1}(0)$, as wanted.

Theorem 4.6 (Formula of Vafa and Intriligator) *(Notations as in Theorem 0.1.) For any $[\omega] \in H^{1,1}(G(k, n))$ the set $C^{[\omega]}$ of critical points of $W^{[\omega]}$ is finite and all of these are non-degenerate. Moreover, for any $F \in \mathbb{C}[X_1, \dots, X_k]$ the following formula for the genus g GRW-invariants of $G(k, n)$ holds:*

$$\langle F \rangle_g^{[\omega]} = (-1)^{n+k(k-1)/2} \sum_{\underline{x} \in C^{[\omega]}} \det \left(\frac{\partial^2 W^{[\omega]}}{\partial X_i \partial X_j} \right) (\underline{x})^{g-1} \cdot F(\underline{x}).$$

◇

Proof. What is left is a check of the normalization. This amounts to calculate the residue of $[\Omega] = (-1)^{n-k} X_k^{n-k}$ with respect to $f^{[\omega]}$. We set $a = (-1)^k e^{[\omega](L)}$. Using standard properties of residues one gets

$$\text{Res}_{f^{[\omega]}} \left((-1)^{n-k} X_k^{n-k} \right) = \frac{(-1)^n}{k!} \text{Res}_{\underline{\lambda}^n - \underline{a}} \left(\prod_{i=1}^k \lambda_i^{n-k} \prod_{i < j} (\lambda_i - \lambda_j)^2 \right),$$

where $\underline{\lambda}^n - \underline{a} = (\lambda_1^n - a, \dots, \lambda_k^n - a)$. Note also that $\prod_{i < j} (\lambda_i - \lambda_j)^2$ is the squared Jacobian of Σ and that $k!$ is the degree of Σ . Since terms of degree less than $k(n-1)$ modulo $(\lambda_i^n - a)$ have vanishing residue (Proposition 4.1) only the term $\prod_{i=1}^k \lambda_i^{k-1}$ from the expansion of $\prod_{i < j} (\lambda_i - \lambda_j)^2$ contributes. The coefficient of this term is $(-1)^{k(k-1)/2} \cdot k!$ as one sees by writing $\prod_{i < j} (\lambda_i - \lambda_j)^2$ as determinant of a product of a Vandermonde-matrix with its transposed. What is left is a multiple of the Jacobian $n^k \prod \lambda_i^{n-1}$ of $\underline{\lambda}^n - \underline{a}$, the residue of which is known to be the degree n^k of $\underline{\lambda}^n - \underline{a}$. Putting everything together we get the claimed normalization. ◇

In concluding let us comment on the connection with the mathematical formulation of this formula given by Bertram, Daskalopoulos and Wentworth [BDW]. They showed that for any Riemann surface Σ the moduli space $\mathcal{M}(d, \Sigma)$ (denoted $\mathcal{M}$ in the sequel) of holomorphic maps $f : \Sigma \rightarrow G(k, n)$, $f_*[\Sigma] = d \cdot [L]$ has the expected dimension provided the degree d is sufficiently large. Moreover, the compactification of $\mathcal{M}$ as Grothendieck quot scheme $\overline{\mathcal{M}}_Q(d, \Sigma)$ (denoted $\overline{\mathcal{M}}$ in the sequel), which parametrizes sheaf quotients $\mathcal{O}_\Sigma^n \rightarrow \mathcal{F} \rightarrow 0$ with fixed Hilbert-polynomial, is generically reduced with irreducible reduction; the universal quotient sheaf $\tilde{\mathcal{F}}$ on $\Sigma \times \mathcal{M}$ has a *locally free* kernel $\tilde{\mathcal{E}}$ extending $\text{ev}^* S$ as subsheaf of $\mathcal{O}_{\Sigma \times \overline{\mathcal{M}}}^n$, S the tautological bundle on $G(k, n)$. In particular it makes sense to talk about the Chern classes $c_i(\tilde{\mathcal{E}})$. Let $\iota : \{t\} \times \overline{\mathcal{M}} \rightarrow \Sigma \times \overline{\mathcal{M}}$ be the inclusion

for some $t \in \Sigma$. Then for $i_1, \dots, i_k$ with $\sum_\nu \nu i_\nu = \dim \mathcal{M} = k(n-k)(1-g) + dn$ they declared

$$\langle X_1^{i_1} \cdots X_k^{i_k} \rangle_\Sigma := \left(\iota^* c_1(\tilde{\mathcal{E}})^{\wedge i_1} \wedge \cdots \wedge \iota^* c_k(\tilde{\mathcal{E}})^{\wedge i_k} \right) [\overline{\mathcal{M}}].$$

Presumably this invariant does not in general coincide with the GRW-invariant: As we tried to explain in Remark 1.2 the GRW-invariants are computed from the Chern classes of $\tilde{\text{ev}}^*(S)$, where $\tilde{\text{ev}} : \Gamma \rightarrow G(k, n)$ is an extension of $\text{ev} : \Sigma \times \mathcal{M} \rightarrow G(k, n)$ to some blow-up $\pi : \Gamma \rightarrow \Sigma \times \overline{\mathcal{M}}$. The trouble is that $\pi^* \tilde{\mathcal{E}}$, though a locally free subsheaf of $\tilde{\text{ev}}^* S$ of the same rank k , it need not coincide with the latter. For instance this always happens if $\tilde{\mathcal{F}}$ has torsion since then $\mathcal{O}_\Gamma^n / \pi^* \tilde{\mathcal{E}} \simeq \pi^* \tilde{\mathcal{F}}$ has torsion as well. Additional contributions thus come precisely from the first k Chern classes of the torsion sheaf $\tilde{\text{ev}}^* \mathcal{O}(S) / \pi^* \tilde{\mathcal{E}}$ (one should desingularize Γ to make sense of these Chern classes). A general vanishing theorem for these seems to be unlikely.

References

- [AsSa] A. Astashkevich and V. Sadov, *Quantum cohomology of Partial Flag Manifolds $F_{n_1 \dots n_k}$* , preprint hep-th 9401130.
- [BDW] A. Bertram, G. Daskalopoulos and R. Wentworth, *Gromov invariants for holomorphic maps from Riemann surfaces to Grassmannians*, preprint, 1993.
- [Be] A. Bertram, *Towards a Schubert calculus for maps from a Riemann surface to a Grassmannian*, preprint alg-geom 9403007.
- [Fi] G. Fischer, *Complex analytic geometry*, LNM 538, Springer, 1976.
- [Fu] W. Fulton, *Intersection Theory*, Springer, 1984.
- [GiKi] A. Givental and B. Kim, *Quantum Cohomology of Flag Manifolds and Toda Lattices*, preprint hep-th 9312096.
- [Gr] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*, Inv. Math. **82** (1985), 307–347.
- [GrHa] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley, 1978.
- [Ha] R. Hartshorne, *Residues and duality*, LNM 20, Springer, 1966.
- [In] K. Intriligator, *Fusion Residues*, Mod. Phys. Lett. A **38** (1991), 3543–3556.
- [Pi] S. Piunikhin, *Quantum and Floer cohomology have the same ring structure*, preprint hep-th 9401130.

- [RRW] M.S. Rafi, J. Rosenthal and X. Wang, *Degree of the generalized Plücker embedding of a quot scheme and quantum cohomology*, preprint hep-th 9402011.
- [Ru] Y. Ruan, *Topological sigma model and Donaldson type invariants in Gromov theory*, Preprint 3/1993, Michigan State University.
- [RuTi] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, in preparation. Announcement to appear in Math. Res. Lett.
- [Ts] A.K. Tsikh, *Multidimensional Residues and Their Applications*, Transl. of Math. Monographs **103**, AMS, 1992.
- [Va1] C. Vafa, *Topological Landau-Ginzburg models*, Mod. Phys. Lett. A **6** (1991), 337–346.
- [Va2] C. Vafa, *Topological Mirrors and Quantum Rings*, in: Essays on Mirror Manifolds (ed. S.-T. Yau), International Press 1992, 96–119.
- [Wi1] E. Witten: *Topological sigma models*, Comm. Math. Phys. **118** (1988), 353–386.
- [Wi2] E. Witten: *On the structure of the topological phase of two-dimensional gravity*, Nucl. Phys. B **340** (1990), 281–332.
- [Wi3] E. Witten: *The Verlinde algebra and the cohomology of the Grassmannian*, preprint hep-th 9312104.